

READ SAFETY DIRECTIONS BEFORE OPENING OR USING



SPREADWET 1000

ACTIVE CONSTITUENT : 1000 g/L ALKOXYLATED ALCOHOLS

**Premium quality *Biodegradable*
wetting and spreading agent
for use with
insecticides, fungicides
and herbicides.**

5 LITRES

Directions for use

Situation	Application Rate	Critical comments
Horticultural sprays Insecticides and Fungicides Knapsack	25ml/100 L 4ml/15 L	Recommended for addition to all horticultural sprays except where formulation has sufficient wetter.
Protected glasshouse horticultural crops	10mL/100Ltr	Rate for sensitive crops in protected glasshouse situation.
Weedkilling sprays Low vol. boom spray Knapsack	100 - 150ml/100 L 4ml/15 L	Refer to individual herbicide labels for specific recommendations.
Glyphosate	100 – 200 mL/100L	
Paraquat and Diquat Boom spray Knapsack	100ml/100 L 15ml/15 L	Under very high dilution rates for maximum efficiency a higher rate of Spreadwet 1000 is used to ensure even coverage of plant foliage and to increase absorption.
Soil Residual Herbicides e.g. Diuron, Simazine, Atrazine	100 - 150ml/100 L	When applying soil active herbicides such as DIURON flowable herbicide to standing weeds add Spreadwet 1000 to thoroughly wet foliage. Check specific product labels for individual recommendations.
Fire fighting Water tanker Knapsack	10ml/100 L 2ml/100 L	Spreadwet 1000 increases the wetting power of water which assists in quick control and mopping up of fires. In cases where water may not be readily available and a less drenching spray is needed a tankful will cover a larger area.

NOT TO BE USED FOR ANY PURPOSE OTHER THAN INDICATED ON THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

GENERAL INSTRUCTIONS

Spreadwet 1000 is a premium quality biodegradable nonionic surfactant for agricultural and horticultural container sprays. It contains 100% w/v active ingredients suitable for use with fungicides, insecticides and herbicides. Spreadwet 1000 reduces the surface tension of spray droplets causing them to spread and adhere to the target in a continuous film. This assists penetration and uptake of the spray on to waxy or hairy plant surfaces, fungous lesions, disease infection sites and insects. Spreadwet 1000 mixes easily with all types of water and will not injure tender foliage if used as directed.

HERBICIDE SPRAYS

Spreadwet 1000 can be added to many herbicide sprays to increase their efficiency. Wetter should be added only where directed on individual labels, as use with selective herbicides outside label directions may reduce crop selectivity. For folia active herbicides spray to the point where leaves and stems are just wet. Spraying beyond this point will encourage run-off and loss of spray chemicals. When spraying hard to wet waxy or hairy leafed plants increased rates of Spreadwet 1000 can increase activity. When treating dense, tall vegetation with folia herbicides it is important to thoroughly wet leaves and stems for efficient herbicide performance.

MIXING

Spreadwet 1000 mixes easily with water and should be added directly to the spray tank after the addition of other components.

PROTECTION OF WILDLIFE, FISH, CRUSTACEA AND THE ENVIRONMENT

Toxic to fish, although rapidly biodegradable. Do not contaminate streams, rivers or waterways with the chemical or used container.

STORAGE AND DISPOSAL

Keep out of reach of Children. Store in closed, original in a well ventilated area, as cool as possible.

Triple or, preferably, pressure rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on-site.

Break, crush, puncture and bury empty containers in a local authority landfill. If not available bury the containers below 500mm in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and roots. Empty containers and product should not be burnt.

SAFETY DIRECTIONS

Avoid contact with skin or eyes. Wear rubber gloves, overalls and goggles or face shield.

FIRST AID

If poisoning occurs contact a doctor or National Poisons Centre 0800 POISON (0800 764 766). If swallowed rinse mouth out with water. Give plenty of water to drink. If vomiting occurs give further water. Seek immediate medical assistance. If skin contact occurs remove contaminated clothing and wash skin thoroughly. If in eyes, hold eyes open and flood with water for at least 15 minutes and see a doctor. Additional information is listed in the Material Safety Data Sheet.

CONDITIONS OF SALE

The use of Spreadwet 1000 being beyond the control of the manufacturer, no warrantee expressed or implied is given by SST NZ LIMITED regarding its suitability, fitness or efficiency for any purpose for which it is used by the buyer, whether in accordance with the directions or not and SST NZ LIMITED accepts no responsibility for any consequences whatsoever resulting from the use of this product.

Batch No.

Date of Manufacture

SST NZ LTD

P.O. Box 6028, Wellesley Street
Level 10, 44 Wellesley Street, Auckland, NEW ZEALAND
PHONE 09 309 8209 Co. No. 1499396

IN AN EMERGENCY PHONE NUMBER IS
NATIONAL POISONS CENTRE
0800 POISON (0800 764 766)