

# New Product



**Analysis:** 2%N, 11%P, 7.5%K + Amino Acids

**Description:** A newly developed N-P-K liquid fertiliser plus biostimulant.









Wuxal AminoPhos contains a plant extract rich in Amino acids, and naturally occurring Auxin. (IAA)

Particularly indicated for application for the improvement and stimulation of root development and the promotion of greater root mass, resulting in the improved overall vigour and health of treated crops.









Can be applied as a direct soil treatment, as a foliar fertiliser, or through fertigation systems.

Due to the nature of the formulation, the nutritional components are highly systemic. Consequently Wuxal AminoPhos is particularly suitable in activating the plants response to stress related conditions.

## **Key benefits & features:**

-  Promotes vigorous and healthy root growth.
-  Nutrients highly plant available.
-  Improves the P and K status of the crop.
-  Improves fruit size and quality.
-  Promotes strong and healthy plant growth.
-  Activates the plants response to stress related conditions.
-  Can be applied through all usual high and low volume spraying equipment.
-  Compatible with a wide range of pesticides.\*

## **Key Crops & Application Rates:**

-  Onions: 5L/Ha. Apply 10-14 days after emergence. Repeat at weekly intervals.
-  Potatoes: 5L/Ha. Apply at tuber initiation then repeat twice at 10-14 day intervals.
-  Brassicas: 5L/Ha. Apply 2-3 weeks following planting out with a maximum of 4 treatments at fortnightly intervals.
-  Lettuce: 3L/Ha. Apply at fortnightly intervals with the first treatment 14 days after planting.
-  Strawberries: (Foliar Sprays) 3L/Ha. 3 applications starting before bloom up until beginning of fruit colouring. (Fertigation: 8-10L/Ha. Repeat applications until start of fruit colouring.
-  Citrus: (Foliar Sprays) 3L/Ha. Apply a maximum of 4 sprays at fortnightly intervals.
-  Ornamental & Cut Flower Crops: 300ml/100L water. Apply as a soil/media drench with 4-5 treatments at fortnightly intervals.
-  Turf Surfaces: (Established) 200-250ml/100m<sup>2</sup>. Repeat at monthly intervals particularly during summer stress.

**Pack Size:** 20L

\* Pesticide compatibility: When mixing with pesticides for the first time, test on a small scale before committing the entire crop!